

# Enzymatic synthesis of poly(alkylene succinate)s: Influence of reaction conditions

Doris Pospiech <sup>1</sup>, Renata Choinska <sup>2</sup>, Daniel Flugrat <sup>1,3</sup>, Karin Sahre <sup>1</sup>, Dieter Jehnichen <sup>1</sup>, Andreas Korwitz <sup>1</sup>, Anett Werner <sup>4</sup>, Brigitte Voit <sup>1,3</sup>

<sup>1</sup> Leibniz-Institut für Polymerforschung Dresden e.V., Hohe Str. 6, 01069 Dresden, Germany; [pospiech@ipfdd.de](mailto:pospiech@ipfdd.de) (D.P.); [karin.sahre@primacom.net](mailto:karin.sahre@primacom.net) (K.S.); [djeh@ipfdd.de](mailto:djeh@ipfdd.de) (D.J.); [korwitz@ipfdd.de](mailto:korwitz@ipfdd.de) (A.K.); [voit@ipfdd.de](mailto:voit@ipfdd.de) (B.V.)

<sup>2</sup> Prof. Waclaw Dąbrowski Institute of Agricultural and Food Biotechnology, Warsaw, Poland; [renata.choinska@ibprs.pl](mailto:renata.choinska@ibprs.pl) (R.C.)

<sup>3</sup> Technische Universität Dresden, Chemistry of Polymers, 01069 Dresden, Germany; [daniel.flugrat@mailbox.tu-dresden.de](mailto:daniel.flugrat@mailbox.tu-dresden.de) (D.F.); [voit@ipfdd.de](mailto:voit@ipfdd.de) (B.V.)


<sup>4</sup> Technische Universität Dresden, Institute of Natural Materials Technology, 01069 Dresden, Germany; [anett.werner@tu-dresden.de](mailto:anett.werner@tu-dresden.de) (A.W.)

\* Correspondence: [pospiech@ipfdd.de](mailto:pospiech@ipfdd.de); Phone: (+49-351-4658497)

## Supporting Information

- 1 GC/MS spectra of the gas phase product evolved in the enzymatic polycondensation of divinyladipate with 1,4-butanediol
- 2 <sup>1</sup>H NMR spectra

1 GC/MS spectra of the gas phase product evolved in the enzymatic polycondensation of divinyladipate with 1,4-butanediol




**Kommentiert [M1]:** This figure is of low resolution. Please provide a sharper one.

**Figure S1.** Headspace GC/MS spectra of products found in the gas phase during polycondensation of divinyladipate with 1,4-butanediol (from top to bottom): 1) gas phase product ( $t = 1.58$  s, cetaldehyde;  $t = 3.68$  s, toluene); 2) empty vial before measurement; 3) empty vial after measurement.

2  $^1\text{H}$  NMR spectra

Assignment of signals for PBA:



$\delta$  ( $\text{CDCl}_3$ ) (ppm): 4.08 (1, 1', m, 4H); 3.68 (t, 2H,  $-\text{CH}_2\text{-OH}$ ); 2.32 (3', m, 4H); 1.69 (2, 2', m, 4H); 1.65 (3, m, 4H); 1.24 ( $-\text{OCO-CH}_3$ , t, 3H) ppm.




Figure S2.  $^1\text{H}$  NMR spectrum of PBA4 (in  $\text{CDCl}_3$ ).




Figure S3.  $^1\text{H}$  NMR spectrum of PBA5-1 (in  $\text{CDCl}_3$ ).




Figure S4.  $^1\text{H}$  NMR spectrum of PBA6 (in  $\text{CDCl}_3$ ).





Figure S5.  $^1\text{H}$  NMR spectrum of PBA2 (in  $\text{CDCl}_3$ ).

Assignment of signals for PBS:



$\delta$  (CDCl<sub>3</sub>) (ppm): 4.10 (1, 1', m, 4H); 3.67 (t, 2H, -CH<sub>2</sub>-OH); 2.60 (3, 3', m, 4H); 1.69 (2, 2', m, 4H) ppm.




Figure S6. <sup>1</sup>H NMR spectrum of PBS1-2 (in CDCl<sub>3</sub>).




Figure S7.  $^1\text{H}$  NMR spectrum of PBS7 (in  $\text{CDCl}_3$ ).




Figure S8.  $^1\text{H}$  NMR spectrum of PBS5 (in  $\text{CDCl}_3$ ).